

First Aid for Seizures



Jennifer

www.onedayicansay.com

Jreyepilepsydiaries

(Complex partial, psychomotor, temporal lobe)

(Convulsive, generalized tonic-clonic, grand mal)

1. Recognize common symptoms



2. Follow first-aid steps



People who've had this type of seizure should be fully conscious and aware before being left on their own. Make sure they know the date, where they are, and where

they're going next. Confusion may last longer than the seizure itself and may be hazardous. If full awareness does not return, call for medical assistance.

EFA 341 | EFA 342
Rev 2/2010



Most seizures in people with epilepsy are not medical emergencies. They end after a minute or two without harm and usually do not require a trip to the emergency room.

But sometimes there are good reasons to call for emergency help. A seizure in someone who does not have epilepsy could be a sign of a serious illness.

Other reasons to call an ambulance include:

- A seizure that lasts more than 5 minutes
- No "epilepsy" or "seizure disorder" identification
- Slow recovery, a second seizure, or difficulty breathing afterwards
- Pregnancy or other medical diagnosis
- Any signs of injury or sickness



ENDEPILEPSY.org

This publication was made possible by a grant from the Centers for Disease Control and Prevention (grant number 5U58DP000606-04) and its contents are solely the responsibility of its authors and do not necessarily represent the official views of the CDC. © 2008, 2010 Epilepsy Foundation of America, Inc.